SYNTHESIS AND ANTICANCER ACTIVITY OF 3-(2,4-DICHLOROPHENOXYMETHYL)-6-ARYL-7H-[1,2,4] TRIAZOLO[3,4-B][1,3,4]THIADIAZINES DERIVATIVES

Demchenko D.A., Yanchenko V.A., Grynevych A.I.

State Centre of Innovation Biotechnologies. 30 Donetska str., Kyiv, Ukraine diana demch@ukr.net

It is known that condensed 4-aminotriazoles exhibit a wide spectrum of biological activity. A range of 7H-[1,2,4]triazolo[3,4-b][1,3,4]thiadiazines 4 was obtained in high overall yields.

C1 O O NHNH₂ C1 O NHNH₂ C1 NH_NH₂ C1 NH_NH₂ C1 NH_NH₂ C1 NH_NH₂ C1 NH_N SH NH₂NH₂ C1
$$R = F$$
, OMe, OCHF₂

Anticancer activity for these compounds was studied at the United States National Cancer Institute (NCI, Bethesda, Maryland, USA). Results obtained show that these compounds are promising for creating of a new drug for the treatment of melanoma on their basis. It was found that anticancer properties of compounds 4 increase with increasing of electron-donor properties of the substituent in the para position. Thus, compound 4 (R = F) inhibits the development of melanoma cells LOX IMVI to 36.73% better than 5-fluorouracil, 4 (R = OCHF₂) - to 80.29%, and 4 (R = OCH₃) - to 90.79%, respectively. This trend is observed in all 8 varieties of cancerous melanoma cells.